

LETTER TO THE EDITOR

Open Access

Answer to comment on “sleep quality, arousal and pain thresholds in migraineurs: a blinded controlled polysomnographic study”

Morten Engstrøm^{1,2*}, Knut Hagen^{1,3}, Marte Bjørk^{4,5} and Trond Sand^{1,2}**Abstract**

We discuss the comments on our article “Sleep quality, arousal and pain thresholds in migraineurs. A blinded controlled polysomnographic study” published in JHP 2013 Feb 14;14(1):12. We hypothesize that migraineurs need more sleep than healthy controls and more sleep than they manage to achieve. Some migraineurs probably have a decreased ability to process incoming stimuli. Increased spontaneous pain may follow either sleep restriction or sleep disturbance. A comparison of migraineurs with attack onset related to sleep, migraineurs with attack onset not related to sleep and controls will be reported in another paper.

Keywords: Migraine; Sleep; Sleep deprivation; Arousal; Pain

Sir,

We thank the group at Institute of Neurology, Catholic University in Rome for the comments [1] on our article [2]. We are delighted to get this opportunity to improve and clarify the interpretation of our findings. It is especially gratifying to answer comments from this group as the publication from 2006 “Dysfunction of arousal system in sleep-related migraine without aura” by Della Marca et al. [3] has been an important and inspiring article during our work with this project.

In our study the migraineurs reported normal sleep times and increased daytime tiredness while we found polysomnographic signs indicating foregoing sleep deprivation. We therefore hypothesized that migraineurs need more sleep than healthy controls and more sleep than they manage to achieve. The relevance of sleep deprivation in migraineurs is amplified by signs of even stronger sleep pressure in the early migraine phase as reduced latency to sleep onset was present.

Both reduced and increased sleep and change in sleep pattern are reported as migraine triggers [4-7]. Even small delays in bedtime can probably release an attack if

the need for sleep is increased. Long sleep is not the most frequently reported migraine trigger and the importance of long sleep itself might be questioned as weekend migraine seems infrequent [8]. To speculate, long sleep periods are probably associated with increased sleep depth which itself seems to be a risk factor for migraine [2,9]. Furthermore, a period of planned sleep restriction most likely is less stress- and harmful than repeated and irregular disturbances during sleep, at least when total sleep time are equal. Repeated disturbances tend to increase light sleep while sleep restriction will increase the more restful slow wave sleep [10]. However, sleep restriction or -deprivation also increase arousal thresholds and thereby also the seriousness of respiratory disturbances [11]. This reasoning is consistent with the notion that arousals are a part of homeostatic adaptive response. High levels of (fast) arousals are probably constitutional or related to repeated occasional disturbances (of variable etiology) during sleep. After a disturbed night the sleep depth and arousal threshold tend to increase [11]. Consequently increased sleep pressure might increase the sleep quality per se but also make one more susceptible to respiratory disturbances and latent sleep apnea. It is therefore difficult to evaluate differences in pain related to sleep disturbance versus sleep restriction. However, there are reports indicating

* Correspondence: morten.engstrom@ntnu.no

¹Department of Clinical Neurosciences, PB 8905, MTF5, Norwegian University of Science and Technology, Trondheim N-7489, Norway

²Department of Neurology and Clinical Neurophysiology, St. Olavs Hospital, Trondheim N-7006, Norway

Full list of author information is available at the end of the article

increased spontaneous pain after sleep restriction [12-14] and also after sleep disturbance [10].

Sleep deprivation also reduce the ability to process incoming stimuli manifested by increased pain sensitivity and consistent with reported increased-, skin-, light noise-, smell sensitivity and testing of pain thresholds among migraineurs [15-21]. However, it is uncertain if affected migraineurs are able to recover and normalize arousal levels whether insomnia is present or not. Reasonable treatment most likely should aim at not making things worse. So bottom line we think some migraineurs have reduced ability to process incoming stimuli, but to us it seems that this reduction is related to a sleep-deprived state. However, another constitutional "sleep-deprived phenotype" with some characteristics similar to narcoleptic patients in sleep [1] is possible.

In our study we did not perform CAP- scoring, but alternatively scored D- and K-bursts similar to CAP A1 bursts [22]. We intend to report results for migraineurs with attack onset related to sleep (sleep migraine) and compare them with migraineurs with attack onset not related to sleep and controls in another submitted paper. We will there also comment on slow bursts, the reported apparent paradoxical finding of reduced arousal- and increased awakening index found in the migraineurs described in our published paper in JHP [2].

Abbreviation

CAP: Cyclic alternating pattern.

Competing interests

The authors have no competing interest to disclose.

Authors' contributions

ME prepared the drafts of this letter. TS has revised several drafts. MB and KH have revised the final version. All authors read and approved the final manuscript.

Author details

¹Department of Clinical Neurosciences, PB 8905, MTF5, Norwegian University of Science and Technology, Trondheim N-7489, Norway. ²Department of Neurology and Clinical Neurophysiology, St. Olavs Hospital, Trondheim N-7006, Norway. ³Norwegian National Headache Centre, St. Olavs Hospital, Trondheim N-7006, Norway. ⁴Department of Neurology, Haukeland University Hospital, Bergen N-5021, Norway. ⁵Department of Clinical Medicine, University of Bergen, Bergen N-5021, Norway.

Received: 29 May 2013 Accepted: 25 June 2013

Published: 1 July 2013

References

- Vollono C, Testani E, Losurdo A, Mazza S, Della Marca G (2013) Migraine, arousal and sleep deprivation: comment on: "sleep quality, arousal and pain thresholds in migraineurs: a blinded controlled polysomnographic study". *J Headache Pain* 14(1):50
- Engstrom M, Hagen K, Bjork MH, Stovner LJ, Gravdahl GB, Stjern M, Sand T (2013) Sleep quality, arousal and pain thresholds in migraineurs: a blinded controlled polysomnographic study. *J Headache Pain* 14(1):12
- Della Marca G, Vollono C, Rubino M, Di Trapani G, Mariotti P, Tonali PA (2006) Dysfunction of arousal systems in sleep-related migraine without aura. *Cephalalgia* 26(7):857-864
- Andress-Rothrock D, King W, Rothrock J (2010) An analysis of migraine triggers in a clinic-based population. *Headache* 50(8):1366-1370
- Hauge AW, Kirchmann M, Olesen J (2011) Characterization of consistent triggers of migraine with aura. *Cephalalgia* 31(4):416-438
- Kelman L (2007) The triggers or precipitants of the acute migraine attack. *Cephalalgia* 27(5):394-402
- Martin PR (2010) Behavioral management of migraine headache triggers: learning to cope with triggers. *Curr Pain Headache Rep* 14(3):221-227
- Alstadhaug KB, Salvesen R, Bekkelund S (2007) Weekend migraine. *Cephalalgia* 27(4):343-346
- Houle TT, Butschek RA, Turner DP, Smitherman TA, Rains JC, Penzien DB (2012) Stress and sleep duration predict headache severity in chronic headache sufferers. *Pain* 153(12):2432-2440
- Smith MT, Edwards RR, McCann UD, Haythornthwaite JA (2007) The effects of sleep deprivation on pain inhibition and spontaneous pain in women. *Sleep* 30(4):494-505
- Bonnet MH, Doghramji K, Roehrs T, Stepanski EJ, Sheldon SH, Walters AS, Wise M, Chesson AL Jr (2007) The scoring of arousal in sleep: reliability, validity, and alternatives. *J Clin Sleep Med* 3(2):133-145
- Haack M, Mullington JM (2005) Sustained sleep restriction reduces emotional and physical well-being. *Pain* 119(1-3):56-64
- Haack M, Lee E, Cohen DA, Mullington JM (2009) Activation of the prostaglandin system in response to sleep loss in healthy humans: potential mediator of increased spontaneous pain. *Pain* 145(1-2):136-141
- Kundermann B, Sperl J, Huber MT, Krieg JC, Lautenbacher S (2004) Sleep deprivation affects thermal pain thresholds but not somatosensory thresholds in healthy volunteers. *Psychosom Med* 66(6):932-937
- Schwedt TJ, Krauss MJ, Frey K, Gereau RW (2011) Episodic and chronic migraineurs are hypersensitive to thermal stimuli between migraine attacks. *Cephalalgia* 31(1):6-12
- Sand T, Zhitniy N, Nilsen KB, Helde G, Hagen K, Stovner LJ (2008) Thermal pain thresholds are decreased in the migraine preattack phase. *Eur J Neurol* 15(11):1199-1205
- Burstein R, Cutrer MF, Yarnitsky D (2000) The development of cutaneous allodynia during a migraine attack clinical evidence for the sequential recruitment of spinal and supraspinal nociceptive neurons in migraine. *Brain* 123(Pt 8):1703-1709
- Zappaterra M, Guerzoni S, Cainazzo MM, Ferrari A, Pini LA (2011) Basal cutaneous pain threshold in headache patients. *J Headache Pain* 12(3):303-310
- Bigal ME, Ashina S, Burstein R, Reed ML, Buse D, Serrano D, Lipton RB, Group A (2008) Prevalence and characteristics of allodynia in headache sufferers: a population study. *Neurology* 70(17):1525-1533
- Lovati C, D'Amico D, Bertora P, Raimondi E, Rosa S, Zardoni M, Bussone G, Mariani C (2010) Correlation between presence of allodynia and sleep quality in migraineurs. *Neurol Sci* 31(Suppl 1):S155-S158
- Olesen J (2004) The international classification of headache disorders: 2nd edition, vol 24 suppl 1. *Cephalalgia* 24:9-160, 2004/02/26 edn
- Terzano MG, Parrino L, Smerieri A, Chervin R, Chokroverty S, Guilleminault C, Hirshkowitz M, Mahowald M, Moldofsky H, Rosa A, Thomas R, Walters A (2002) Atlas, rules, and recording techniques for the scoring of cyclic alternating pattern (CAP) in human sleep. *Sleep Med* 3(2):187-199

doi:10.1186/1129-2377-14-56

Cite this article as: Engstrøm et al.: Answer to comment on "sleep quality, arousal and pain thresholds in migraineurs: a blinded controlled polysomnographic study". *The Journal of Headache and Pain* 2013 **14**:56.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com